

Perfusion Index Measurement in Preterm Newborns during the First Days of Life

Bianchi A., Ossola S., Stifani A., Bossi A., Agosti M. *Pediatric Academic Societies Annual Meeting* 2012:3853.669

Background

Perfusion Index (PI) is a non-invasive, continuous indicator of peripheral perfusion and it seems to be a marker of severe conditions from different causes. Reference values of PI in term newborns have been recently published, but only few data exist about PI values in preterm newborns. *Objective:* To determine PI values in the first days of life in preterm newborns, even in correlation with gestational age (GA) and birth weight (BW).

Design/Methods

We studied 35 neonates born ≤ 32 weeks GA without malformation syndromes (mean GA: $29,33 \pm 2,04$ weeks; mean BW: 1192 ± 395 g). The newborns were divided according to GA in 24-28 weeks (group1) and 29-32 weeks (group2) and to BW ≤ 1000 g (group A) and >1000 g (group B). PI was measured at 25-48, 49-72, 73-96 hours of life; it was recorded continuously for 10' (average of values recorded every 20") at the right hand and foot by Masimo Radical-7 pulse oximeter.

Results

Median PI at 25-48 hours of life is 1,30, at 49-72 hours is 1,52 and at 73-96 hours is 1,45. At 25-48, 49-72, 73-96 hours group1 has lower PI than group2 and group A has significantly lower PI than group B.

Conclusion

Median PI is lower at 25-48 hours of life, when preterm neonates are at higher risk of peripheral hypoperfusion. PI grows in the following days reflecting the hemodynamic changes.

PI has a positive correlation with GA and a statistical significance with BW at 25-48 hours and at 73-96 hours. PI measurement is an effective method to monitor preterm newborns in the first days of life.

age	Group 1	Group 2	p	Group A	Group B	p
25-48	1,28 $\pm$ 0,28	1,46 $\pm$ 0,53	ns	1,17 $\pm$ 0,29	1,52 $\pm$ 0,49	<0,05
49-72	1,48 $\pm$ 0,62	1,69 $\pm$ 0,57	ns	1,51 $\pm$ 0,49	1,66 $\pm$ 0,64	ns
73-96	1,53 $\pm$ 0,70	1,71 $\pm$ 0,59	ns	1,36 $\pm$ 0,75	1,75 $\pm$ 0,56	<0,05

